MULTIMEDIA UNIVERSITY

STUDENT ID NO

MULTIMEDIA UNIVERSITY

FINAL EXAMINATION

TRIMESTER 1, 2022/2023 (TERM ID: 2215)

BQT1614 – QUANTITATIVE ANALYSIS

(All Sections/Groups)

20 FEBRUARY 2023
9.00 am – 12.00 pm
(3 Hours)

INSTRUCTIONS TO STUDENT

1. This question paper consists of **TWELVE (12)** pages (excluding the cover page).

2. Answer **ALL** questions. The marks distributions are given in parentheses.

3. The Formulae Sheet and the Statistical Tables are provided.

QUESTION 1 [20 MARKS]

A sample of 30 students was picked at random. The obtained information pertains to their weekly coffee expenditures (in RM) and was documented as follows:

21	22.5	23	25.5	26	26	27	27.5	28	28
29	31.5	32	33	33	33	35	35.6	36	37
37	37	38	38.5	39	40	42.5	45	51.5	56

a) Calculate the mean and median of the data.

[4 marks]

b) Calculate the standard deviation and coefficient of variation of the data.

[4 marks]

c) Find the 98% confidence interval for the proportion of customers that order more than 35.

[5 marks]

d) If there were another 5 customers that may overspend : 56, 56.7, 57.5 and 59 and 60 were added to the data, how will the answer in (a), (b) and the dispersion of the data changed?

[7 marks]

Continued...

QUESTION 2 [20 MARKS]

a) A survey of 500 people was made to determine the preferred way for people to order fast food. The results are summarized in the following table:

Preferred way to order	**Gender**		**Total**
	Male	*Female*	
Dine Inside	16	44	60
Order inside to go	97	53	150
Order at the drive-through	140	150	290
Total	253	247	500

i. If a respondent is selected at random, what is the probability that he or she prefers to dine inside?

[3 marks]

ii. If a male in the survey is selected, find the probability that he prefers to order at the drive-through?

[3 marks]

iii. Determine whether the events gender and the preferred way for people to order fast food are independent by providing statistical evidence.

[4 marks]

b) In 2022, it is claimed that 60% of Malaysians would be influenced by social influencer marketing to purchase any product they see. If a sample of 10 Malaysian adult is selected;

i. what is the probability that 5 will purchase once they see social influencer marketing advertises

[2 marks]

ii. what is the mean and standard deviation that one will purchase once they see social influencer marketing advertises?

[2 marks]

Continued...

c) The points earned on the Weekly Quiz in the subject of Statistics, comprising 30 points, are nearly normally distributed, with a mean of 24 and a standard deviation of 6.

i. What is the probability that a person will scores between 20 and 30 points?

[3 marks]

ii. What is the minimum points does a person have to score to be in the highest 20% of scores for the Weekly Quiz in the subject of Statistics?

[3 marks]

QUESTION 3 [20 MARKS]

a) Santo tyre manufacturing company produces two types of tires, A and B. Out of random sample of 500 type A tires, 60 tires are defective and out of a random sample of 720 type B tires, 85 are defective. Is there a significant difference between proportions of defective of type A and type B tires? Use $\alpha = 0.05$.

[12 marks]

b) In an audit firm previously the mean salary of the auditors is RM 6500 per month. A random sample of 8 auditors has a mean salary of RM7100 per month. The standard deviation of the sample is RM300 per month. At the 0.01 level of significance, can we conclude that the mean salary of the auditors has changed?

[8 marks]

QUESTION 4 [20 MARKS]

Lucas-Copiers, sells and services the copying machine. As part of its standard service contract, the company agrees to perform routine service on this copier. To obtain information about the time it takes to perform routine service, Lucas-copiers has collected data for 11 service calls. The data are as follows:

Number of Copiers Serviced	Number of Minutes Required
4	109
2	58
5	138
7	189
1	37
3	82
4	103
5	134
2	68
4	112
6	154

Continued...

a) Identify the independent and dependent variables.

[2 marks]

b) Assuming a linear relationship, use the least-squares method to obtain the regression line equation.

[6 marks]

c) Interpret the meaning of the slope obtained in (b).

[2 marks]

d) Use the regression line obtained in (b) to predict the time required to service 9 copiers. Is the estimation reliable?

[4 marks]

e) Determine the coefficient of correlation and interpret its meaning.

[3 marks]

f) Determine the coefficient of determination and interpret its meaning.

[3 marks]

QUESTION 5 [20 MARKS]

a) Sam made a down payment of RM20,000 toward the purchase of a car. He secured a 9- year loan from a bank at 3% per year compounded monthly. Under the term of the loan agreement, he has to make a payment of RM715 at the end of each month.
   i. What is the purchase price of the car? [4 marks]
   ii. What is the interest paid? [2 marks]

b) A newly married couple has $15,000 toward the purchase of a house. For the type of house they are interested in buying they estimate that a $20,000 down payment will be necessary. How long will the money have to be invested at 10% compounded quarterly to grow to $20,000? [3 marks]

c) A company establishes a fund to upgrade a plant in 5 years at an estimated cost of $1,500,000. How much should be invested each quarter into an account paying 9% compounded quarterly? How much interest will the account earn in the 5 years?

[5 marks]

Continued...

d) The table below shows the prices and quantities of several items sell in their Co-op Shop in a private school.

Item	2017		2022	
	Price (RM)	Quantity	Price (RM)	Quantity
Set of Pen	5	120	7	140
Set of Writing Book	30	24	32	28
Uniforms	220	20	230	18
Shoes	180	25	195	26

i. Determine the Laspeyres price index for 2022 by using 2017 as the base year.
[3 marks]

ii. Determine the Paasche quantity index for 2022 by using 2017 as the base year.
[3 marks]

End of Question.

APPENDIXES:

I. STATISTICAL FORMULAE

A. DESRIPTIVE STATISTICS

Sample mean: $\bar{X} = \frac{\sum x}{n}$

Sample standard deviation: $S = \sqrt{\frac{\sum x^2}{n-1} - \frac{(\sum x)^2}{n(n-1)}}$

Coefficient of variation (sample): $CV = \frac{S}{\bar{X}} \times 100\%$

Pearson coefficient of skewness: $S_K = \frac{3(\bar{X} - Median)}{S}$

B. PROBABILITY

$P(A \cup B) = P(A) + P(B) - P(A \cap B)$

$P(A \cap B) = P(A) \times P(B)$ if A and B are independent

$P(A \mid B) = P(A \cap B) \div P(B)$

Poisson Probability Distribution

If X follows a Poisson Distribution P (λ) where $P(X = x) = \frac{e^{-\lambda}\lambda^x}{x!}$

then the mean = E(X) = λ and variance = VAR(X) = λ

Binomial Probability Distribution

If X follows a Binomial Distribution B(n, p) where $P(X = x) = {}^nC_x p^x q^{n-x}$

then the mean = E(X) = np and variance = VAR(X) = $=npq$ where $q = 1 - p$

Normal Distribution

If X follows a Normal distribution N(μ , σ) where E(X) = μ and VAR(X) = σ^2

then $Z = \frac{X - \mu}{\sigma}$

C. CONFIDENCE INTERVAL ESTIMATION AN SAMPLE SIZE DETERMINATION

$(100-\alpha)\%$ Confidence Interval for Population Mean:

If σ known, μ: $\bar{x} \pm Z_{\alpha/2}\frac{\sigma}{\sqrt{n}}$

If σ unknown, μ: $\bar{x} \pm t_{\alpha/2,n-1}\frac{s}{\sqrt{n}}$

$(100-\alpha)\%$ Confidence Interval for Population Proportion, π: $p \pm Z_{\alpha/2}\sqrt{\frac{p(1-p)}{n}}$

Sample Size Determination for Population Mean: $n = \left(\frac{(Z_{\alpha/2})\ \sigma}{e}\right)^2$

Sample Size Determination for Population Proportion: $n = \frac{(Z_{\alpha/2})^2 \pi(1-\pi)}{e^2}$

Continued...

D. HYPOTHESIS TESTING

One Sample Mean Test

Z-test: $Z = \dfrac{\bar{x} - \mu}{\dfrac{\sigma}{\sqrt{n}}}$ **t-test:** $t = \dfrac{\bar{x} - \mu}{\dfrac{s}{\sqrt{n}}}$

One Sample Proportion Test

$$Z = \frac{p - \pi}{\sqrt{\dfrac{\pi(1-\pi)}{n}}}$$

Two Sample Mean Test

Z-test: $Z = \dfrac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\dfrac{\sigma_1^2}{n_1} + \dfrac{\sigma_2^2}{n_2}}}$

t-test: $t = \dfrac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{s_p\sqrt{\left(\dfrac{1}{n_1} + \dfrac{1}{n_2}\right)}}$ where $s_p = \sqrt{\dfrac{(n_1-1)s_1^2 + (n_2-1)s_2^2}{n_1 + n_2 - 2}}$

Two Sample Proportion Test

$$Z = \frac{(p_1 - p_2) - (\pi_1 - \pi_2)}{\sqrt{\bar{p}(1-\bar{p})\left[\dfrac{1}{n_1} + \dfrac{1}{n_2}\right]}} \quad \text{where} \quad \bar{p} = \frac{x_1 + x_2}{n_1 + n_2}; \quad p_1 = \frac{x_1}{n_1}; \quad p_2 = \frac{x_2}{n_2}$$

E. REGRESSION ANALYSIS

Simple Linear Regression

$$y_i = b_0 + b_1 x_i \text{ where } b_0 = \bar{Y} - b_1\bar{X} \text{ and } b_1 = \frac{\sum XY - \left[\dfrac{\sum X \sum Y}{n}\right]}{\left[\sum X^2 - \left(\dfrac{(\sum X)^2}{n}\right)\right]}$$

Correlation Coefficient

$$r = \frac{n\sum xy - \sum x \sum y}{\sqrt{[n\sum x^2 - (\sum x)^2][n\sum y^2 - (\sum y)^2]}}$$

ANOVA Table for Regression

Source	Degrees of Freedom	Sum of Squares	Mean Squares
Regression	*1*	SSR	MSR = SSR/1
Error/Residual	*n – 2*	SSE	MSE = SSE/(*n – 2*)
Total	*n – 1*	SST	

Test Statistic for Significance of the Predictor Variable

$t_i = \dfrac{b_i}{S_{b_i}}$ and the critical value = $\pm t_{\alpha/2,(n-p-1)}$ where p = number of predictor

Continued.....

F. INDEX NUMBERS

Simple Price Index $P = \frac{p_t}{p_0} \times 100$	**Simple Aggregate Price Index** $P = \frac{\sum p_t}{\sum p_0} \times 100$
Laspeyres Quantity Index $P = \frac{\sum p_0 q_t}{\sum p_0 q_0} \times 100$	**Paasche Quantity Index** $P = \frac{\sum p_t q_t}{\sum p_t q_0} \times 100$
Laspeyres Price Index $P = \frac{\sum p_t q_0}{\sum p_0 q_0} \times 100$	**Paasche Price Index** $P = \frac{\sum p_t q_t}{\sum p_0 q_t} \times 100$
Fisher's Ideal Price Index $\sqrt{(Laspeyres\ Price\ Index)(Paasche\ Price\ Index)}$	**Value Index** $V = \frac{\sum p_t q_t}{\sum p_0 q_0} \times 100$

G. FINANCIAL MATHEMATICS

Simple Interest

Interest: $I = \Pr t$

Accumulated amount: $A = P(1+rt)$

Compounded Interest

Accumulated amount: $A = P(1+i)^n$

Present Value: $P = A(1+i)^{-n}$

Interest: $I = P(1+i)^n - P$

Effective rate of interest

$$r_e = \left(1 + \frac{r}{m}\right)^m - 1$$

Sinking Fund $R = \left[\frac{Si}{(1+i)^n - 1}\right]$	**Amortization** $R = \left[\frac{Pi}{1-(1+i)^{-n}}\right]$
Future Value of an Annuity $A = R\left[\frac{(1+i)^n - 1}{i}\right]$	**Present Value of an Annuity** $P = R\left[\frac{1-(1+i)^{-n}}{i}\right]$

Continued...

II. STATISTICAL TABLE

Table 1

The Upper Tail Area Under the Standard Normal Curve



Z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	0.5000	0.4960	0.4920	0.4880	0.4840	0.4801	0.4761	0.4721	0.4681	0.4641
0.1	0.4602	0.4562	0.4522	0.4483	0.4443	0.4404	0.4364	0.4325	0.4286	0.4247
0.2	0.4207	0.4168	0.4129	0.4090	0.4052	0.4013	0.3974	0.3936	0.3897	0.3859
0.3	0.3821	0.3783	0.3745	0.3707	0.3669	0.3632	0.3594	0.3557	0.3520	0.3483
0.4	0.3446	0.3409	0.3372	0.3336	0.3300	0.3264	0.3228	0.3192	0.3156	0.3121
0.5	0.3085	0.3050	0.3015	0.2981	0.2946	0.2912	0.2877	0.2843	0.2810	0.2776
0.6	0.2743	0.2709	0.2676	0.2643	0.2611	0.2578	0.2546	0.2514	0.2483	0.2451
0.7	0.2420	0.2389	0.2358	0.2327	0.2296	0.2266	0.2236	0.2206	0.2177	0.2148
0.8	0.2119	0.2090	0.2061	0.2033	0.2005	0.1977	0.1949	0.1922	0.1894	0.1867
0.9	0.1841	0.1814	0.1788	0.1762	0.1736	0.1711	0.1685	0.1660	0.1635	0.1611
1.0	0.1587	0.1562	0.1539	0.1515	0.1492	0.1469	0.1446	0.1423	0.1401	0.1379
1.1	0.1357	0.1335	0.1314	0.1292	0.1271	0.1251	0.1230	0.1210	0.1190	0.1170
1.2	0.1151	0.1131	0.1112	0.1093	0.1075	0.1056	0.1038	0.1020	0.1003	0.0985
1.3	0.0968	0.0951	0.0934	0.0918	0.0901	0.0885	0.0869	0.0853	0.0838	0.0823
1.4	0.0808	0.0793	0.0778	0.0764	0.0749	0.0735	0.0721	0.0708	0.0694	0.0681
1.5	0.0668	0.0655	0.0643	0.0630	0.0618	0.0606	0.0594	0.0582	0.0571	0.0559
1.6	0.0548	0.0537	0.0526	0.0516	0.0505	0.0495	0.0485	0.0475	0.0465	0.0455
1.7	0.0446	0.0436	0.0427	0.0418	0.0409	0.0401	0.0392	0.0384	0.0375	0.0367
1.8	0.0359	0.0351	0.0344	0.0336	0.0329	0.0322	0.0314	0.0307	0.0301	0.0294
1.9	0.0287	0.0281	0.0274	0.0268	0.0262	0.0256	0.0250	0.0244	0.0239	0.0233
2.0	0.0228	0.0222	0.0217	0.0212	0.0207	0.0202	0.0197	0.0192	0.0188	0.0183
2.1	0.0179	0.0174	0.0170	0.0166	0.0162	0.0158	0.0154	0.0150	0.0146	0.0143
2.2	0.0139	0.0136	0.0132	0.0129	0.0125	0.0122	0.0119	0.0116	0.0113	0.0110
2.3	0.0107	0.0104	0.0102	0.0099	0.0096	0.0094	0.0091	0.0089	0.0087	0.0084
2.4	0.0082	0.0080	0.0078	0.0075	0.0073	0.0071	0.0069	0.0068	0.0066	0.0064
2.5	0.0062	0.0060	0.0059	0.0057	0.0055	0.0054	0.0052	0.0051	0.0049	0.0048
2.6	0.0047	0.0045	0.0044	0.0043	0.0041	0.0040	0.0039	0.0038	0.0037	0.0036
2.7	0.0035	0.0034	0.0033	0.0032	0.0031	0.0030	0.0029	0.0028	0.0027	0.0026
2.8	0.0026	0.0025	0.0024	0.0023	0.0023	0.0022	0.0021	0.0021	0.0020	0.0019
2.9	0.0019	0.0018	0.0018	0.0017	0.0016	0.0016	0.0015	0.0015	0.0014	0.0014
3.0	0.0013	0.0013	0.0013	0.0012	0.0012	0.0011	0.0011	0.0011	0.0010	0.0010
3.1	0.0010	0.0009	0.0009	0.0009	0.0008	0.0008	0.0008	0.0008	0.0007	0.0007
3.2	0.0007	0.0007	0.0006	0.0006	0.0006	0.0006	0.0006	0.0005	0.0005	0.0005
3.3	0.0005	0.0005	0.0005	0.0004	0.0004	0.0004	0.0004	0.0004	0.0004	0.0003
3.4	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0002
3.5	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002
3.6	0.0002	0.0002	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001
3.7	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001
3.8	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001
3.9	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
4.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000

Continued...

TABLE 4. THE NORMAL DISTRIBUTION FUNCTION

The function tabulated is $\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x} e^{-\frac{1}{2}t^2} dt$. $\Phi(x)$ is the probability that a random variable, normally distributed with zero mean and unit variance, will be less than or equal to x. When $x < 0$ use $\Phi(x) = 1 - \Phi(-x)$, as the normal distribution with zero mean and unit variance is symmetric about zero.



x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$
0·00	0·5000	0·40	0·6554	0·80	0·7881	1·20	0·8849	1·60	0·9452	2·00	0·97725
·01	·5040	·41	·6591	·81	·7910	·21	·8869	·61	·9463	·01	·97778
·02	·5080	·42	·6628	·82	·7939	·22	·8888	·62	·9474	·02	·97831
·03	·5120	·43	·6664	·83	·7967	·23	·8907	·63	·9484	·03	·97882
·04	·5160	·44	·6700	·84	·7995	·24	·8925	·64	·9495	·04	·97932
0·05	0·5199	0·45	0·6736	0·85	0·8023	1·25	0·8944	1·65	0·9505	2·05	0·97982
·06	·5239	·46	·6772	·86	·8051	·26	·8962	·66	·9515	·06	·98030
·07	·5279	·47	·6808	·87	·8078	·27	·8980	·67	·9525	·07	·98077
·08	·5319	·48	·6844	·88	·8106	·28	·8997	·68	·9535	·08	·98124
·09	·5359	·49	·6879	·89	·8133	·29	·9015	·69	·9545	·09	·98169
0·10	0·5398	0·50	0·6915	0·90	0·8159	1·30	0·9032	1·70	0·9554	2·10	0·98214
·11	·5438	·51	·6950	·91	·8186	·31	·9049	·71	·9564	·11	·98257
·12	·5478	·52	·6985	·92	·8212	·32	·9066	·72	·9573	·12	·98300
·13	·5517	·53	·7019	·93	·8238	·33	·9082	·73	·9582	·13	·98341
·14	·5557	·54	·7054	·94	·8264	·34	·9099	·74	·9591	·14	·98382
0·15	0·5596	0·55	0·7088	0·95	0·8289	1·35	0·9115	1·75	0·9599	2·15	0·98422
·16	·5636	·56	·7123	·96	·8315	·36	·9131	·76	·9608	·16	·98461
·17	·5675	·57	·7157	·97	·8340	·37	·9147	·77	·9616	·17	·98500
·18	·5714	·58	·7190	·98	·8365	·38	·9162	·78	·9625	·18	·98537
·19	·5753	·59	·7224	·99	·8389	·39	·9177	·79	·9633	·19	·98574
0·20	0·5793	0·60	0·7257	1·00	0·8413	1·40	0·9192	1·80	0·9641	2·20	0·98610
·21	·5832	·61	·7291	·01	·8438	·41	·9207	·81	·9649	·21	·98645
·22	·5871	·62	·7324	·02	·8461	·42	·9222	·82	·9656	·22	·98679
·23	·5910	·63	·7357	·03	·8485	·43	·9236	·83	·9664	·23	·98713
·24	·5948	·64	·7389	·04	·8508	·44	·9251	·84	·9671	·24	·98745
0·25	0·5987	0·65	0·7422	1·05	0·8531	1·45	0·9265	1·85	0·9678	2·25	0·98778
·26	·6026	·66	·7454	·06	·8554	·46	·9279	·86	·9686	·26	·98809
·27	·6064	·67	·7486	·07	·8577	·47	·9292	·87	·9693	·27	·98840
·28	·6103	·68	·7517	·08	·8599	·48	·9306	·88	·9699	·28	·98870
·29	·6141	·69	·7549	·09	·8621	·49	·9319	·89	·9706	·29	·98899
0·30	0·6179	0·70	0·7580	1·10	0·8643	1·50	0·9332	1·90	0·9713	2·30	0·98928
·31	·6217	·71	·7611	·11	·8665	·51	·9345	·91	·9719	·31	·98956
·32	·6255	·72	·7642	·12	·8686	·52	·9357	·92	·9726	·32	·98983
·33	·6293	·73	·7673	·13	·8708	·53	·9370	·93	·9732	·33	·99010
·34	·6331	·74	·7704	·14	·8729	·54	·9382	·94	·9738	·34	·99036
0·35	0·6368	0·75	0·7734	1·15	0·8749	1·55	0·9394	1·95	0·9744	2·35	0·99061
·36	·6406	·76	·7764	·16	·8770	·56	·9406	·96	·9750	·36	·99086
·37	·6443	·77	·7794	·17	·8790	·57	·9418	·97	·9756	·37	·99111
·38	·6480	·78	·7823	·18	·8810	·58	·9429	·98	·9761	·38	·99134
·39	·6517	·79	·7852	·19	·8830	·59	·9441	·99	·9767	·39	·99158
0·40	0·6554	0·80	0·7881	1·20	0·8849	1·60	0·9452	2·00	0·9772	2·40	0·99180

Continued...

Continued...

TABLE 4. THE NORMAL DISTRIBUTION FUNCTION

x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$
2·40	0·99180	2·55	0·99461	2·70	0·99653	2·85	0·99781	3·00	0·99865	3·15	0·99918
·41	·99202	·56	·99477	·71	·99664	·86	·99788	·01	·99869	·16	·99921
·42	·99224	·57	·99492	·72	·99674	·87	·99795	·02	·99874	·17	·99924
·43	·99245	·58	·99506	·73	·99683	·88	·99801	·03	·99878	·18	·99926
·44	·99266	·59	·99520	·74	·99693	·89	·99807	·04	·99882	·19	·99929
2·45	0·99286	2·60	0·99534	2·75	0·99702	2·90	0·99813	3·05	0·99886	3·20	0·99931
·46	·99305	·61	·99547	·76	·99711	·91	·99819	·06	·99889	·21	·99934
·47	·99324	·62	·99560	·77	·99720	·92	·99825	·07	·99893	·22	·99936
·48	·99343	·63	·99573	·78	·99728	·93	·99831	·08	·99896	·23	·99938
·49	·99361	·64	·99585	·79	·99736	·94	·99836	·09	·99900	·24	·99940
2·50	0·99379	2·65	0·99598	2·80	0·99744	2·95	0·99841	3·10	0·99903	3·25	0·99942
·51	·99396	·66	·99609	·81	·99752	·96	·99846	·11	·99906	·26	·99944
·52	·99413	·67	·99621	·82	·99760	·97	·99851	·12	·99910	·27	·99946
·53	·99430	·68	·99632	·83	·99767	·98	·99856	·13	·99913	·28	·99948
·54	·99446	·69	·99643	·84	·99774	·99	·99861	·14	·99916	·29	·99950
2·55	0·99461	2·70	0·99653	2·85	0·99781	3·00	0·99865	3·15	0·99918	3·30	0·99952

The critical table below gives on the left the range of values of x for which $\Phi(x)$ takes the value on the right, correct to the last figure given; in critical cases, take the upper of the two values of $\Phi(x)$ indicated.

3·075	0·9990	3·263	0·9994	3·731	0·99990	3·916	0·99995
3·105	0·9991	3·320	0·9995	3·759	0·99991	3·976	0·99996
3·138	0·9992	3·389	0·9996	3·791	0·99992	4·055	0·99997
3·174	0·9993	3·480	0·9997	3·826	0·99993	4·173	0·99998
3·215	0·9994	3·615	0·9998	3·867	0·99994	4·417	0·99999
			0·9999		0·99995		1·00000

When $x > 3{\cdot}3$ the formula $1-\Phi(x) \doteqdot \frac{e^{-\frac{1}{2}x^2}}{x\sqrt{2\pi}}\left[1-\frac{1}{x^2}+\frac{3}{x^4}-\frac{15}{x^6}+\frac{105}{x^8}\right]$ is very accurate, with relative error less than $945/x^{10}$.

TABLE 5. PERCENTAGE POINTS OF THE NORMAL DISTRIBUTION

This table gives percentage points $x(P)$ defined by the equation

$$\frac{P}{100} = \frac{1}{\sqrt{2\pi}}\int_{x(P)}^{\infty} e^{-\frac{1}{2}t^2}\,dt.$$

If X is a variable, normally distributed with zero mean and unit variance, $P/100$ is the probability that $X \geqslant x(P)$. The lower P per cent points are given by symmetry as $-x(P)$, and the probability that $|X| \geqslant x(P)$ is $2P/100$.



P	$x(P)$	P	$x(P)$	P	$x(P)$	P	$x(P)$	P	$x(P)$	P	$x(P)$
50	0·0000	5·0	1·6449	3·0	1·8808	2·0	2·0537	1·0	2·3263	0·10	3·0902
45	0·1257	4·8	1·6646	2·9	1·8957	1·9	2·0749	0·9	2·3656	0·09	3·1214
40	0·2533	4·6	1·6849	2·8	1·9110	1·8	2·0969	0·8	2·4089	0·08	3·1559
35	0·3853	4·4	1·7060	2·7	1·9268	1·7	2·1201	0·7	2·4573	0·07	3·1947
30	0·5244	4·2	1·7279	2·6	1·9431	1·6	2·1444	0·6	2·5121	0·06	3·2389
25	0·6745	4·0	1·7507	2·5	1·9600	1·5	2·1701	0·5	2·5758	0·05	3·2905
20	0·8416	3·8	1·7744	2·4	1·9774	1·4	2·1973	0·4	2·6521	0·01	3·7190
15	1·0364	3·6	1·7991	2·3	1·9954	1·3	2·2262	0·3	2·7478	0·005	3·8906
10	1·2816	3·4	1·8250	2·2	2·0141	1·2	2·2571	0·2	2·8782	0·001	4·2649
5	1·6449	3·2	1·8522	2·1	2·0335	1·1	2·2904	0·1	3·0902	0·0005	4·4172

Continued...

TABLE 10. PERCENTAGE POINTS OF THE t-DISTRIBUTION

This table gives percentage points $t_\nu(P)$ defined by the equation

$$\frac{P}{100} = \frac{1}{\sqrt{\nu\pi}} \frac{\Gamma(\frac{1}{2}\nu+\frac{1}{2})}{\Gamma(\frac{1}{2}\nu)} \int_{t_\nu(P)}^{\infty} \frac{dt}{(1+t^2/\nu)^{\frac{1}{2}(\nu+1)}}.$$

Let X_1 and X_2 be independent random variables having a normal distribution with zero mean and unit variance and a χ^2-distribution with ν degrees of freedom respectively; then $t = X_1/\sqrt{X_2/\nu}$ has Student's t-distribution with ν degrees of freedom, and the probability that $t \geqslant t_\nu(P)$ is $P/100$. The lower percentage points are given by symmetry as $-t_\nu(P)$, and the probability that $|t| \geqslant t_\nu(P)$ is $2P/100$.



The limiting distribution of t as ν tends to infinity is the normal distribution with zero mean and unit variance. When ν is large interpolation in ν should be harmonic.

P	40	30	25	20	15	10	5	2·5	1	0·5	0·1	0·05
$\nu = 1$	0·3249	0·7265	1·0000	1·3764	1·963	3·078	6·314	12·71	31·82	63·66	318·3	636·6
2	0·2887	0·6172	0·8165	1·0607	1·386	1·886	2·920	4·303	6·965	9·925	22·33	31·60
3	0·2767	0·5844	0·7649	0·9785	1·250	1·638	2·353	3·182	4·541	5·841	10·21	12·92
4	0·2707	0·5686	0·7407	0·9410	1·190	1·533	2·132	2·776	3·747	4·604	7·173	8·610
5	0·2672	0·5594	0·7267	0·9195	1·156	1·476	2·015	2·571	3·365	4·032	5·893	6·869
6	0·2648	0·5534	0·7176	0·9057	1·134	1·440	1·943	2·447	3·143	3·707	5·208	5·959
7	0·2632	0·5491	0·7111	0·8960	1·119	1·415	1·895	2·365	2·998	3·499	4·785	5·408
8	0·2619	0·5459	0·7064	0·8889	1·108	1·397	1·860	2·306	2·896	3·355	4·50	5·041
9	0·2610	0·5435	0·7027	0·8834	1·100	1·383	1·833	2·262	2·821	3·250	4·297	4·781
10	0·2602	0·5415	0·6998	0·8791	1·093	1·372	1·812	2·228	2·764	3·169	4·144	4·587
11	0·2596	0·5399	0·6974	0·8755	1·088	1·363	1·796	2·201	2·718	3·106	4·025	4·437
12	0·2590	0·5386	0·6955	0·8726	1·083	1·356	1·782	2·179	2·681	3·055	3·930	4·318
13	0·2586	0·5375	0·6938	0·8702	1·079	1·350	1·771	2·160	2·650	3·012	3·852	4·221
14	0·2582	0·5366	0·6924	0·8681	1·076	1·345	1·761	2·145	2·624	2·977	3·787	4·140
15	0·2579	0·5357	0·6912	0·8662	1·074	1·341	1·753	2·131	2·602	2·947	3·733	4·073
16	0·2576	0·5350	0·6901	0·8647	1·071	1·337	1·746	2·120	2·583	2·921	3·686	4·015
17	0·2573	0·5344	0·6892	0·8633	1·069	1·333	1·740	2·110	2·567	2·898	3·646	3·965
18	0·2571	0·5338	0·6884	0·8620	1·067	1·330	1·734	2·101	2·552	2·878	3·610	3·922
19	0·2569	0·5333	0·6876	0·8610	1·066	1·328	1·729	2·093	2·539	2·861	3·579	3·883
20	0·2567	0·5329	0·6870	0·8600	1·064	1·325	1·725	2·086	2·528	2·845	3·552	3·850
21	0·2566	0·5325	0·6864	0·8591	1·063	1·323	1·721	2·080	2·518	2·831	3·527	3·819
22	0·2564	0·5321	0·6858	0·8583	1·061	1·321	1·717	2·074	2·508	2·819	3·505	3·792
23	0·2563	0·5317	0·6853	0·8575	1·060	1·319	1·714	2·069	2·500	2·807	3·485	3·768
24	0·2562	0·5314	0·6848	0·8569	1·059	1·318	1·711	2·064	2·492	2·797	3·467	3·745
25	0·2561	0·5312	0·6844	0·8562	1·058	1·316	1·708	2·060	2·485	2·787	3·450	3·725
26	0·2560	0·5309	0·6840	0·8557	1·058	1·315	1·706	2·056	2·479	2·779	3·435	3·707
27	0·2559	0·5306	0·6837	0·8551	1·057	1·314	1·703	2·052	2·473	2·771	3·421	3·690
28	0·2558	0·5304	0·6834	0·8546	1·056	1·313	1·701	2·048	2·467	2·763	3·408	3·674
29	0·2557	0·5302	0·6830	0·8542	1·055	1·311	1·699	2·045	2·462	2·756	3·396	3·659
30	0·2556	0·5300	0·6828	0·8538	1·055	1·310	1·697	2·042	2·457	2·750	3·385	3·646
32	0·2555	0·5297	0·6822	0·8530	1·054	1·309	1·694	2·037	2·449	2·738	3·365	3·622
34	0·2553	0·5294	0·6818	0·8523	1·052	1·307	1·691	2·032	2·441	2·728	3·348	3·601
36	0·2552	0·5291	0·6814	0·8517	1·052	1·306	1·688	2·028	2·434	2·719	3·333	3·582
38	0·2551	0·5288	0·6810	0·8512	1·051	1·304	1·686	2·024	2·429	2·712	3·319	3·566
40	0·2550	0·5286	0·6807	0·8507	1·050	1·303	1·684	2·021	2·423	2·704	3·307	3·551
50	0·2547	0·5278	0·6794	0·8489	1·047	1·299	1·676	2·009	2·403	2·678	3·261	3·496
60	0·2545	0·5272	0·6786	0·8477	1·045	1·296	1·671	2·000	2·390	2·660	3·232	3·460
120	0·2539	0·5258	0·6765	0·8446	1·041	1·289	1·658	1·980	2·358	2·617	3·160	3·373
∞	0·2533	0·5244	0·6745	0·8416	1·036	1·282	1·645	1·960	2·326	2·576	3·090	3·291

End of Page